

Erythromycin Stearate Tablet USP

Product Information :

Pharmaceutical form : Tablet

Category : Prescription Medicine

Pack : 1x10/Tabs

Brand name : Erythromycin Stearate

Content : Erythromycin (500mg)

Therapeutic Use : Treatment of Bacterial infections



Product Description :

What is Erythromycin Stearate Tablet USP?

Erythromycin Stearate Tablet USP is used in the treatment of bacterial infections of the respiratory tract, such as bronchitis, pneumonia, Legionnaires' disease (lung infection), and pertussis (whooping cough; an infection that can lead to serious coughing); sexually transmitted infections, such as syphilis; and ear, intestine, gynaecological, urinary tract, and skin problems. It's also used to keep rheumatic fever from returning. Erythromycin belongs to the macrolide antibiotics family of drugs. It works by preventing bacteria from growing.

How to Use

- Erythromycin is available in the form of a tablet, capsule, delayed release tablet and an oral solution (liquid) to take by mouth.
- It's commonly taken each six hours (4 times a day), every eight hours (3 times a day), or every 12 hours (2 times a day), both with and without food.
- Do not chew or crush the capsules or tablets; swallow them entirely with a full glass of

water.

- Even though you feel fine, keep taking Erythromycin Stearate Tablet USP. Stopping erythromycin without consulting your doctor is not a good option.
- Erythromycin is also used to protect against heart infections in persons who are having dentistry or other operations done. Consider the risks of using this medicine for your illness with your doctor.

Precaution - Erythromycin Stearate Tablet USP

- If you are allergic to erythromycin, any other drugs, or any of the substances in erythromycin capsules, tablets, or suspension, consult your doctor.
- If you're receiving astemizole, cisapride, dihydroergotamine, ergotamine, pimozone, or terfenadine, inform your doctor. If you are using one or more of these medications, your doctor will probably advise you not to take erythromycin.
- Tell your doctor about any prescription and OTC medications, nutraceuticals, and herbal products you're using or planning to use. Many other medicines may interfere with erythromycin, so be sure to notify about all of your current medications. Your doctor will need to adjust your prescription dosages or keep a close eye on you for side effects.
- If you have or have ever had a prolonged QT interval (a heart condition that can cause fainting or abnormal heartbeat), a rapid heartbeat, weak magnesium or potassium levels in the blood, or liver disease, notify your doctor.
- If you have a pregnancy, planning to become pregnant, or breastfeeding, notify your doctor. Call your doctor when you become pregnant while using Erythromycin Stearate Tablet USP.
- Tell your dentist that you are receiving erythromycin if you are getting dental surgery.

Benefits of Erythromycin Tablet

Erythromycin Tablet is a type of antibiotic medication that can be used in the treatment of a variety of bacterial illnesses. Ear, nose, throat, lungs, and skin infections are among them. It stops the infection-causing germs from growing and relieves the infection. Use it for as long as your doctor suggests and don't miss any doses. This will ensure that all germs are eliminated and that resistance does not develop.

Side Effects of Erythromycin

The majorities of side effects are minor and will go away as your body adapts to the medication. If they don't go away or you're concerned about them, see your doctor.

- Nausea, vomiting, diarrhoea, stomach pain, and appetite loss are all possible side effects. This drug may help to alleviate these symptoms if taken with food. Tell your doctor immediately if any of these side effects persist or worsen.
- Remember that your doctor provided Erythromycin Stearate Tablet USP because he or she believes the advantage to you exceeds the risk of adverse effects. The majority of people who take this medicine do not have any substantial adverse effects.
- If you have any severe side effects, such as persistent vomiting, yellowing eyes or skin, black urine, serious stomach/abdominal pain, tiredness, muscle weakness, problems speaking, impaired vision, drooping eyelids, or hearing loss, contact your doctor immediately.
- If you have any very significant adverse effects, such as extreme dizziness, fainting, or an irregular heartbeat, seek medical attention immediately.

Tips For Using Erythromycin Tablet

- Erythromycin Stearate Tablet USP has been given by your doctor to treat your infection and relieve your symptoms.
- Even though you feel better, do not miss any doses and complete the entire course of treatment. Stopping it too soon may cause the infection to return and become more difficult to cure.
- It should be taken 60 minutes before or 2 hours after eating.
- Antacids should not be used two hours before or after consuming Erythromycin tablets.
- Diarrhoea may develop as a side effect, but it should subside once your treatment is over. If it doesn't go away or if you notice blood in your stool, contact your doctor.
- If you experience an itching rash, swelling of the face, throat, or tongue, or difficulty breathing while consuming the Erythromycin Tablet, stop taking it and contact your doctor right away.

Erythromycin Stearate Tablet USP Supplier

Healthiza is one of the leading Erythromycin Stearate Tablet USP supplier and manufacturer, offering the best quality medicine globally through a validated cold chain secure shipment process. We use this process to manufacture, store and deliver medicines at a very specific temperature. Customers can avail of this premium quality pharma product at an affordable price from us.



GET IN TOUCH HEALTHIZA LIFESCIENCE PVT LTD

EMAIL : info@healthizalife.com | sales@healthizalife.com

PHONE : +91- 93289 40307

ADDRESS : A-907, Signature II, Sarkhej 382210 Ahmedabad, India.

WEBSITE : www.healthizalife.com