## **Polidocanol Injection**

### **Product Information:**

Pharmaceutical form: Injection

Category : Anti Cancer

Pack: 1x1vial

Brand name : Polidocanol Content : POLIDOCANOL 2ml

Therapeutic Use: Treatment of varicose veins



## **Product Description:**

## What is Polidocanol Injection?

Polidocanol injection is a prescription medicine used in the treatment with varicose veins. It is used to treat spider veins (varicose veins with a diameter of less than or equal to 1 mm) and simple reticular veins (varicose veins with a diameter of 1 to 3 mm) in the lower limb. Polidocanol is also used to treat large auxiliary saphenous veins that are incompetent, saphenous veins and apparent varicose veins in the vast saphenous vein system above and below the knee.

#### **How It Works**

Polidocanol is a type of sclerosing agent medication. It affects the endothelium of vascular system locally, causing platelets to clump together and form blood clots, as well as scar tissue, which limits the lumen of some larger veins.

#### **Polidocanol Injection - Side Effects**

The most of side effects of Polidocanol injection are minor and will go away as your body responds to the medication. If they don't go away or you're concerned, see your doctor.

- Itching
- Skin Irritation
- Reactions At The Site Of The Injection
- Hematoma, A Collection Of Blood Outside Of The Blood Vessels
- Hives
- Angioedema (Type Of Allergic Reaction)
- Confusion
- Headache
- Fainting
- Dizziness
- Fever
- Pain
- Trouble Breathing

**Some Tips for Polidocanol** 

- This Polidocanol injection is injected directly into the varicose veins, and the number of injections required is determined by the number of varicose veins present. You will be carefully monitored for any negative side effects while you are being injected.
- To avoid clot formation, you should use a compression bandage or stocking for 2 to 3 days following the injection and for 2 to 3 weeks during the day.
- Along with using the compression bandage, you should walk for 15-20 minutes after therapy.
- For the first 2-3 days after therapy, avoid heavy or intense exertion. Sitting for long periods of time should also be prevented (in a car or airplane).
- After treatment, stay out of the sun for at least 2-3 days.
- Before using ice or heating pads to the treated limb, follow your doctor's instructions.

FAQ

- Does Polidocanol injection cause an increase in heart rate or a drop in heart rate?
- Polidocanol can help to lower your heart rate. It does not, however, raise the heart rate.
- Is Polidocanol associated with weight gain or a rise in blood pressure?
- It raises Blood pressure. However, there has been no clinically significant weight loss associated with the administration of Polidocanol.
- Is Polidocanol injection good for dry eyes?
- Yes, Polidocanol aids in the relief of dry eyes by boosting tear production.
- Is Polidocanol associated with blurred vision or hair loss?
- Blurred vision is a side effect of Polidocanol injection. However, it does not result in hair

loss.

## What effect does Polidocanol have on heart rate?

■ It lowers heart rate via boosting the impact of acetylcholine, a neurotransmitter.

#### **Polidocanol Injection Supplier**

Healthiza is one of the leading Pilocarpine injection supplier and manufacturer, offering the best quality medicine globally through a validated cold chain secure shipment process. We use this process to manufacture, store and deliver medicines at a very specific temperature. Customers can avail of this premium quality pharma product at an affordable price from us.



# GET IN TOUCH HEALTHIZA LIFESCIENCE PVT LTD

**EMAIL**: info@healthizalife.com | sales@healthizalife.com

**PHONE:** +91- 93289 40307

ADDRESS: A-907, Signature II, Sarkhej 382210 Ahmedabad, India.

WEBSITE: www.healthizalife.com