# **Ledipasvir Tablet**

### **Product Information:**

Pharmaceutical form: Tablet

Category : Anti Cancer

Pack: 1x28/Tabs

**Brand name: Ledipasvir Tablet** 

Content: Ledipasvir (90mg) + Sofosbuvir

(400mg)

Therapeutic Use: Treatment of Chronic hepatitis

C virus (HCV) infection



# **Product Description:**

## What is Ledipasvir Tablet?

The ledipasvir tablet is used alone or with a combination of other drugs such as

#### sofosbuvir

, and ribavirin to treat some types of chronic Hepatitis C Virus (a continuous viral infection that destroys the liver) In adults and children 3 years of age and older. Ledipasvir tablet, belong to a class of antiviral drugs known as HCV NS5A inhibitors. It works by preventing the hepatitis C virus from spreading inside the body.

How Ledipasvir Tablet - How It Works

Ledipasvir tablet is an antiviral medication that acts in combination with Sofosbuvir. They act by reducing the amount of Hepatitis C Virus in the body and gradually eliminating the infection from the blood.

Ledipasvir Tablets - How To Use

- This medicine is normally given by mouth once a day, with or without food. Every day, around the same time, take the ledipasvir tablet. Follow the advice on your medicine label carefully.
- Even if you felt Fine, keep taking the ledipasvir tablet. The length of your treatment (eight to twenty-four weeks) is determined by your situation, how well you react to the drug, and whether you have severe adverse effects. Do not discontinue taking ledipasvir without first consulting your expert.

#### **Precaution For Ledipasvir Tablet**

- If you are sensitive to ledipasvir, any other drugs, or any of the substances in the ledipasvir tablet, inform your doctor right away.
- Tell your doctor what other prescription and nonprescription medications, vitamins, and nutritional supplements you are taking or plan to take. Be sure to mention any of the following: medications for HIV taken together such as

## <u>efavirenz</u>

#### emtricitabine

and

#### tenofovir

; elvitegravir, cobicistat, emtricitabine, and, atazanavir and ritonavir, certain medications that suppress the immune system; rifabutin; rifampin; rifapentine;

#### rosuvastatin

; simeprevir;

#### sofosbuvir

; and warfarin. Your doctor may tell you not to take ledipasvir if you are taking one or more of these medications or may monitor you carefully for side effects. Many other medications may also interact with the ledipasvir tablet.

- Tell your doctor about any herbal supplements you're taking while taking the ledipasvir tablet.
- If you're using cimetidine, ranitidine, famotidine, or nizatidine for indigestion, heartburn, or ulcers, notify your doctor. You may be told to take these 12 hours before or after the ledipasvir tablet by your doctor.
- If you've had a liver transplant or have the human immunodeficiency virus (HIV), tell your doctor. Also, if you have or have had had any form of liver illness other than hepatitis C, renal disease, or are on dialysis, tell your doctor.

#### **Ledipasvir Tablets - Side Effects**

The majority of side effects are minor and will go away as your body adapts to the medication. If they don't go away or you're concerned about them, see your doctor.

- Tiredness, headaches, nausea, diarrhoea, and difficulties sleeping are all possible side effects. Consult your doctor right away if any of these side effects persist or worsen.
- Remember that your doctor provided this medication because he or she believes the advantage to you exceeds the risk of adverse effects. The majority of people who take this medicine do not have any substantial adverse effects.
- This medicine rarely causes a severe allergic reaction. However, if you detect any symptoms of a major allergic response, such as a rash, itching/swelling, extreme dizziness, or problems breathing, seek medical treatment right once.
- Contact your doctor if you have any other side effects not listed above.

#### Some Tips for Ledipasvir Tablet

- Ledipasvir tablet is a drug that is used to treat hepatitis c Virus infection.
- Take it with food and at the same time every day if possible.
- You should take the medicine every day for 12 to 24 weeks.
- When medicine dosages have been skipped the likelihood of treatment failure rises.
  Make sure you take all of your medications at the same time.
- Fatigue, dizziness, and vision problems are all possible side effects. Don't drive or do

anything else that requires attention until you have a better understanding of how it affects you.

- Headaches are a possible side effect of ledipasvir. Drink lots of fluids and take a pain reliever that suits your needs. If it doesn't go away, consult your doctor.
- Your liver activity and the quantity of the hepatitis C virus in your body may be monitored by your doctor on a regular basis.
- Do not avoid or skip this medicine without your doctor's direction.

#### **Ledipasvir Tablet Supplier**

Healthiza is one of the leading Ledipasvir Tablet supplier and manufacturer, offering the best quality medicine globally through a validated cold chain secure shipment process. We use this process to manufacture, store and deliver medicines at a very specific temperature. Customers can avail of this premium quality pharma product at an affordable price from us.



# GET IN TOUCH HEALTHIZA LIFESCIENCE PVT LTD

**EMAIL**: info@healthizalife.com | sales@healthizalife.com

**PHONE:** +91- 93289 40307

ADDRESS: A-907, Signature II, Sarkhej 382210 Ahmedabad, India.

WEBSITE: www.healthizalife.com