

Nelfinavir Mesylate Tablet

Product Information :

Pharmaceutical form : Tablet

Category : Anti Cancer

Pack : 1x100/Tabs

Brand name : Nelfinavir Mesylate Tablet

Content : Nelfinavir mesylate 250mg

Therapeutic Use : To treat human immunodeficiency virus (HIV) infection



Product Description :

What is Nelfinavir Mesylate Tablet?

Nelfinavir Mesylate Tablet is a medicine that is used to treat HIV infection in combination with other drugs. Nelfinavir belongs to a class of drugs known as protease inhibitors. It works by lowering HIV levels in the bloodstream. It can reduce your risk of acquiring AIDS and HIV-related diseases such as serious infections or cancer.

Nelfinavir Mesylate Tablet - How To Use

- Nelfinavir is available as a tablet and a powder for oral use. It's commonly taken with food 2 or 3 times a day. Every day, around the same time, take the Nelfinavir Mesylate Tablet. Follow the advice on your medicine label completely.
- If you can't eat it, you can put the tablet in a glass and melt it with a tiny amount of water. Drink the liquid right after thoroughly mixing it.
- Nelfinavir can be mixed with water, milk, formula, soy milk, or other nutritional supplements. To get the entire dose, mix well and consume all of the liquid straight away.

- Nelfinavir Mesylate Tablet is an antiretroviral drug that prevents HIV infection. Even if you feel fine, keep taking nelfinavir. Do not discontinue taking nelfinavir without first consulting your physician.

How Nelfinavir Mesylate Tablet Works

Nelfinavir is an antiretroviral drug. It operates by interacting with a protein (protease) that HIV-infected cells need to produce new viruses. This prevents the spread of new viruses.

Nelfinavir Mesylate Tablet - Precautions

Before taking Nelfinavir

- If you are sensitive to nelfinavir, any other medicines, or any of the components in Nelfinavir Mesylate Tablet, inform your doctor right away.
- If you're taking bromocriptine, dihydroergotamine, ergoloid mesylates, ergotamine, or methylergonovine, lovastatin, lurasidone, midazolam, pimozide, quinidine, rifampin,

[sildenafil](#)

, simvastatin, If you are consuming one or more of these medications, your doctor will probably advise you not to take nelfinavir.

- Oral contraceptives may be less effective if you take nelfinavir. While using this medicine, you should utilise a different method of birth control.
- Tell your doctor about any prescription and over-the-counter drugs, vitamins, and nutraceuticals you're taking or planning to take. Make a point of mentioning one or more of the following: anticoagulants ('blood thinners') like warfarin;

[azithromycin](#)

; bosentan; certain calcium-channel blocking medications like

[amlodipine](#)

, felodipine, isradipine, nimodipine, and nisoldipine; carbamazepine; certain cholesterol-lowering medications like

atorvastatin

and rosuvastatin; colchicine; delavirdine; fluticasone Your doctor may need to adjust your medication dosages carefully.

- Do not take didanosine for more than 2 hours before or after taking nelfinavir.
- If you have diabetes, haemophilia or liver illness, tell your doctor.
- If you are pregnant, planning to become pregnant, or breastfeeding, notify your doctor. If you have HIV and are taking nelfinavir, you should not breastfeed.
- Tell your doctor that you are receiving nelfinavir if you are getting surgery, especially dental surgery.
- You should be advised that your body fat may accumulate or migrate to other parts of your body, including your upper back, neck, breasts, and stomach.
- Even if you do not have diabetes, you should be aware that you may have hyperglycemia (blood sugar spikes) when taking this medicine.
- You should be aware that when taking

HIV drugs

, your immune system may become stronger and start to fight other infections already present in your body. It's possible that you'll experience symptoms of those ailments as a result of this. If you notice any new or worsening complications during your nelfinavir treatment, notify your doctor right once.

Side Effects of Nelfinavir Mesylate Tablet

- The majorities of side effects are minor and will go away as your body adapts to the medication. If they don't go away or you're concerned about them, see your doctor. Some of the side effects are given
- Rash, headache, dizziness, and vomiting are all symptoms of the flu. Urine with crystals, Nausea, Pain in the abdomen, Indigestion, Throat ache, Increased levels of liver enzymes Diarrhea, Urine with blood, Cough, Changes in taste, Hypercholesterolemia is a condition in which a person's cholesterol (high cholesterol), Increased blood triglyceride

levels, Urinary protein.

Nelfinavir Mesylate Tablet - Some Tips For use

- It's best to take a Nelfinavir Mesylate Tablet with food.
- Even if you feel better, do not miss any doses and complete the entire course of treatment.
- It can make you feel dizzy or sleepy. Do not drive or engage in any activity that needs concentration until you have a better understanding of how it affects you.
- Your Doctor will examine your liver function on a regular basis. If you suffer abdominal pain, a loss of appetite, darker urine, or yellowing of the eyes or skin, call your doctor (jaundice).
- If you are pregnant, planning a pregnancy, or breastfeeding, tell your doctor.

Nelfinavir Mesylate Tablet Supplier

Healthiza is one of the leading Nelfinavir Mesylate Tablet supplier and manufacturer, offering the best quality medicine globally through a validated cold chain secure shipment process. We use this process to manufacture, store and deliver medicines at a very specific temperature. Customers can avail of this premium quality pharma product at an affordable price from us.



GET IN TOUCH HEALTHIZA LIFESCIENCE PVT LTD

EMAIL : info@healthizalife.com | sales@healthizalife.com

PHONE : +91- 93289 40307

ADDRESS : A-907, Signature II, Sarkhej 382210 Ahmedabad, India.

WEBSITE : www.healthizalife.com