

Pregabalin Tablet 75 mg

Product Information :

Pharmaceutical form : Tablet

Category : Prescription Medicine

Pack : 1x15

Brand name : Pregabalin

Content : Pregabalin Tablet 75 mg

Therapeutic Use : Treatment of Neuropathic pain, Epilepsy/Seizures & Fibromyalgia



Product Description :

What is Pregabalin Tablet 75 mg?

Pregabalin Tablet 75 mg is used in the treatment of neuropathic pain (pain caused by damaged nerves) that can occur in your arms, hands, fingers, legs, feet, or toes if you have diabetes and postherpetic neuralgia. This medication together with other drugs is used in the treatment of some types of seizures in adults and children aged one month and up.

Pregabalin belongs to the anticonvulsant drug class. It works by reducing the number of pain impulses sent out by the body's injured nerves.

How to Use Pregabalin

- Pregabalin is available as a capsule, liquid, or tablet for oral administration. Pregabalin tablets and oral solution are commonly taken 2 to 3 times a day, both with and without food. Pregabalin Tablet 75 mg is typically taken once a day, after dinner.
- Do not cut, chew, or crush the tablets; they must be swallowed whole.
- During the first week of treatment, your doctor will likely begin you on smaller dosages of

pregabalin and gradually raise it.

- It should be taken exactly as prescribed. Pregabalin has the potential to become addictive. Do not take a higher dose, take it more frequently, or longer than recommended.
- Pregabalin's full benefit may take a few weeks or longer to emerge. Even though you feel fine, keep taking it. Even if you have side effects such as unexpected changes in behaviour or mood, do not discontinue taking pregabalin without consulting your doctor.
- You may get withdrawal symptoms such as difficulty falling or staying asleep, nausea, diarrhoea, migraines, or seizures if you stop taking it suddenly. Your doctor will most likely reduce your dose slowly over the course of at least one week.

Precaution - Pregabalin Tablet 75 mg

- If you have an allergy to pregabalin, any other drugs, or any of the components in pregabalin preparations, tell your doctor.
- Tell your doctor about any additional drugs, vitamins, nutritional supplements, or herbal products you're taking or planning to use. Your doctor will need to adjust your prescription dosages or keep a close eye on you for adverse effects.
- If you consume or have ever drunk a lot of alcohol, tell your doctor. If you have or have ever had swelling of the eyes, face, lips, tongue, or throat; vision issues; heart problems; bleeding problems or a low amount of platelets in your blood; or lung, heart, or kidney disease, notify your doctor.
- Tell your doctor or dentist that you are receiving Pregabalin Tablet 75 mg if you are getting surgery, including dental surgery.
- Talk to your doctor if you are pregnant or plan to become pregnant. If you're breastfeeding, let your doctor know. Call your doctor when you become pregnant while taking pregabalin.
- Pregabalin has the potential of making you dizzy or sleepy. Do not drive a car, operate machinery, or engage in other highly unsafe tasks until you have a better understanding of how this medicine affects you.

How it Works

Pregabalin is a type of antiepileptic drug. When used to treat epilepsy, it is thought to prevent seizures by lowering inappropriate brain waves. It's also used to treat nerve pain by interrupting pain signals that flow through injured nerves and into the brain. It is thought to function for anxiety treatment by inhibiting the production of particular chemical transmitters (neurotransmitters) that cause anxiety.

Side Effects of Pregabalin Tablet

The majorities of side effects are minor and will go away as your body adapts to the medication. If they don't go away or you're worried about them, see your doctor.

- It's possible that you'll have drowsiness, dizziness, headaches, dry mouth, nausea, diarrhoea, and weight gain. Tell your doctor right away if any of these side effects persist or worsen.
- Remember that your doctor recommended Pregabalin Tablet 75 mg because he or she believes the advantage to you exceeds the risk of adverse effects. Many persons who use this medicine have no major adverse effects.
- If you experience any serious side effects, such as impaired vision, abnormal bleeding/bruising, muscle pain/tenderness/weakness (particularly with fever or unusual weariness), or swollen hands/ankles/feet, contact your doctor immediately.
- If you or a family member/caregiver detect any unusual changes in your mood, thoughts, or behaviour, such as indicators of depression, or ideas about harming yourself, contact your doctor immediately.

Some Tips for Using Pregabalin

- Pregabalin should be used according to your doctor's instructions for dosage and duration.
- It could make you sleepy. Do not drive a vehicle or do anything else that requires attention until you have a better understanding of how it affects you.
- It may cause eyesight blurring or loss. If you detect any changes in your vision, contact your doctor immediately.

- If you experience any unexpected changes in mood or behaviour, contact your doctor.
- Stopping the drug without first consulting your doctor could make your underlying illness worse.

Pregabalin Tablet 75 mg Supplier

Healthiza is one of the leading Pregabalin Tablet 75 mg supplier and manufacturer, offering the best quality medicine globally through a validated cold chain secure shipment process. We use this process to manufacture, store and deliver medicines at a very specific temperature. Customers can avail of this premium quality pharma product at an affordable price from us.



GET IN TOUCH HEALTHIZA LIFESCIENCE PVT LTD

EMAIL : info@healthizalife.com | sales@healthizalife.com

PHONE : +91- 93289 40307

ADDRESS : A-907, Signature II, Sarkhej 382210 Ahmedabad, India.

WEBSITE : www.healthizalife.com