

Biperidin 2 mg Tablet

Product Information :

Pharmaceutical form : Tablet

Category : Prescription Medicine

Pack : 1x10

Brand name : Biperidin

Content : Biperidin 2mg Tablets

Therapeutic Use : Anti parkinson



Product Description :

What is Biperidin 2 mg Tablet?

Biperidin 2 mg Tablet is a prescription medicine used in the treatment of Parkinson's disease alone or in combination with other medications. As the disease effects fade, this drug improves muscular control and reduces stiffness, allowing for more typical body motions.

Biperidin is also used to treat severe muscle reactions and other adverse effects related with certain drugs used to treat severe nausea or neurological, mental, or emotional problems (e.g., chlorpromazine, perphenazine, prochlorperazine, promethazine, thioridazine).

Biperidin belongs to the Anticholinergic drug class.

How to use Biperidin Tablet

Follow your doctor's instructions when taking Biperidin. If you don't understand something, ask your doctor to explain it to you.

- Each dose should be taken with a full glass of water.
- If Biperidin disturbs your stomach, take it after a meal.

- Biperidin 2 mg Tablet can be used one or more times per day. A maximum of eight tablets of Biperidin should be used in a single day (16 mg). Pay attention to your doctor's directions.
- Biperidin should be stored at room temperature, free from moisture and heat.

Side Effects of Biperidin

The majorities of side effects are minor and will go away as your body adapts to the medication. If they don't go away or you're worried, consult your doctor. Avoid taking Biperidin 2 mg Tablet and get emergency medical help if you experience any symptoms of significant adverse effects:

- An allergic reaction (trouble breathing; throat closure; swelling of the lips, tongue, or face; or hives);
- Unusually high fever;
- Abnormal heartbeat
- Anxiety, hallucinations, disorientation, agitation, hyperactivity, or loss of consciousness are some of the symptoms that can occur.
- Seizures
- Eye Pain
- Rash

Other side effects may also occur. Continue to take Biperidin and consult your doctor if you have any minor adverse effects.

Biperidin 2 mg Tablet Supplier



GET IN TOUCH HEALTHIZA LIFESCIENCE PVT LTD

EMAIL : info@healthizalife.com | sales@healthizalife.com

PHONE : +91- 93289 40307

ADDRESS : A-907, Signature II, Sarkhej 382210 Ahmedabad, India.

WEBSITE : www.healthizalife.com